arXiv:1103.3109v1 [math.GN] 16 Mar 2011

On γ-Semi-Continuous Functions

Sabir *Hussain*[1], Bashir *Ahmad*[2] and Takashi *Noiri*[3]
[1]Department of Mathematics, Islamia University Bahawalpur, Pakistan.
Present Address: Department of Mathematics, Yanbu University,
P. O. Box 31387, Yanbu, Saudi Arabia.
E. mail: sabiriub@yahoo.com.
[2]Department of Mathematics, King Abdul Aziz University, P. O. Box 80203,
Jeddah 21589, Saudi Arabia.
E. mail: drbashir9@gmail.com.
[3]2949-1 Shiokita-cha, Hinagu, Yatsushiro-shi, Kumamoto-ken, 869-5142, Japan.
E. mail: t.noiri@nifty.com.

Abstract. In this paper, we continue studying the properties of γ-semi-continuous and γ-semi-open functions introduced in [5].

Keywords. γ-closed (open), γ-closure , γ^*-semi-closed (open), γ^*-semi-closure, γ-regular, γ^*-semi-interior, γ-semi-continuous function, γ-semi-open (closed) function.
AMS(2000)Subject Classification. Primary 54A05, 54A10, 54D10.

1 Introduction

N. Levine [11] introduced the notion of semi-open sets in topological spaces. A. Csaszar [7,8] defined generalized open sets in generalized topological spaces. In 1975, Maheshwari and Prasad [12] introduced concepts of semi-T_1-spaces and semi-R_0-spaces. In 1979, S. Kasahara [10] defined an operation α on topological spaces. Carpintero, et. al [6] introduced the notion of α-semi-open sets as a generalization of semi-open sets. B. Ahmad and F.U. Rehman [1, 14] introduced the notions of γ-interior, γ-boundary and γ-exterior points in topological spaces. They also studied properties and characterizations of (γ ,β)-continuous mappings introduced by H. Ogata [13]. In [2-4], B. Ahmad and S. Hussain introduced the concept of γ_0-compact, γ^*-regular, γ-normal spaces and explored their many interesting properties. They initiated and discussed the concepts of γ^*-semi-open sets , γ^*-semi-closed sets, γ^*-semi-closure, γ^*-semi-interior points in topological spaces [5,9]. In [9], they introduced Λ_s^γ-set and Λ^{s^γ}-set by using γ^*-semi-open sets. Moreover, they also introduced γ-semi-continuous function and γ-semi-open (closed) functions in topological spaces and established several interesting properties.

In this paper, we continue studying the properties of γ-semi-continuous functions and γ-semi-open function introduced by B. Ahmad and S. Hussain [5].

Hereafter, we shall write space in place of topological space in the sequel.

2 Preliminaries

We recall some definitions and results used in this paper to make it self-contained.

Definition [13]. Let (X,τ) be a space. An operation $\gamma : \tau \rightarrow$ P(X) is a function from τ to the power set of X such that V $\subseteq V^\gamma$, for each V $\in \tau$, where V^γ denotes the value of γ at V. The operations defined by γ(G) = G, γ(G) = cl(G) and γ(G) = intcl(G) are examples of operation γ.

Definition [13]. Let A be a subset of a space X. A point x $\in$ A is said to be γ-interior point of A, if there exists an open nbd N of x such that $N^\gamma \subseteq$ A. The set of all such points is denoted by int_γ(A). Thus

$$int_\gamma \text{ (A)} = \{x \in A : x \in N \in \tau \text{ and } N^\gamma \subseteq A\} \subseteq A.$$

Note that A is γ-open [13] iff A =int_γ(A). A set A is called γ- closed [13] iff X-A is γ-open.

Definition [10]. A point x $\in$ X is called a γ-closure point of A $\subseteq$ X, if $U^\gamma \cap A \neq \phi$, for each open nbd U of x. The set of all γ-closure points of A is called γ-closure of A and is denoted by cl_γ(A). A subset A of X is called γ-closed, if $cl_\gamma(A)$ $\subseteq$A. Note that cl_γ(A) is contained in every γ-closed superset of A.

Definition [14]. The γ-exterior of A, written $ext_\gamma(A)$ is defined as the γ-interior of $(X-A)$. That is, $int_\gamma(A) = ext_\gamma(X-A)$.

Definition [14]. The γ-boundary of A, written $bd_\gamma(A)$ is defined as the set of points which do not belong to γ-interior or the γ-exterior of A.

Definition [13]. An operation γ on τ is said be regular, if for any open nbds U,V of x ∈ X, there exists an open nbd W of x such that $U^{\gamma} \cap V^{\gamma} \supseteq W^{\gamma}$.

Definition [13]. An operation γ on τ is said to be open, if for every open nbd U of each $x \in X$, there exists γ-open set B such that $x \in B$ and $U^{\gamma} \subseteq B$.

Definition [2]. Let $A \subseteq X$. A point $x \in X$ is said to be γ-limit point of A, if $U \cap \{A - \{x\}\} \neq \phi$, where U is a γ-open set in X. The set of all γ-limit points of A denoted A^{d}_{γ} is called γ-derived set.

Definition [9]. A subset A of a space (X,τ) is said to be a γ^{*}-semi-open set, if there exists a γ -open set O such that $O \subseteq A \subseteq cl_{\gamma}(O)$. The set of all γ^{*}-semi-open sets is denoted by $SO_{\gamma^{*}}(X)$.

Definition [5]. A function $f : (X, \tau) \to (Y, \tau)$ is said to be γ-semi-continuous, if for any γ-open B of Y, $f^{-1}(B)$ is γ^{*}-semi-open in X.

Definition [5]. A function $f : (X, \tau) \to (Y, \tau)$ is said to be γ-semi-open (closed), if for each γ-open (closed) set U in X, $f(U)$ is γ^{*}-semi-open (closed) in Y.

Definition [5]. A set A in a space X is said to be γ^{*}-semi-closed, if there exists a γ-closed set F such that $int_{\gamma}(F) \subseteq A \subseteq F$.

Proposition [5]. A subset A of X is γ^{*}-semi-closed if and only if $X - A$ is γ^{*}-semi-open.

Definition 2.1. A subset A of X is said to be γ-semi-nbd of a point $x \in X$, if there exists a γ^{*}-semi-open set U such that $x \in U \subseteq A$.

Definition [9]. Let A be a subset of space X . The intersection of all γ^{*}-semi-closed sets containing A is called γ^{*}-semi-closure of A and is denoted by $scl_{\gamma^{*}}(A)$. Note that A is γ^{*}-semi-closed if and only if $scl_{\gamma^{*}}(A) = A$.

Definition [5]. Let A be a subset of a space X . The union of all γ^{*}-semi-open sets of X contained in A is called γ^{*}-semi-interior of A and is denoted by $sint_{\gamma^{*}}(A)$.

Lemma 2.2. Let A be a subset of a space X. Then $x \in scl_{\gamma^{*}}(A)$ if and only if for any γ-semi-nbd N_x of x in X, $A \cap N_x \neq \phi$.

Proof. Let $x \in scl_{\gamma^{*}}(A)$ .Suppose on the contrary, there exists a γ-semi-nbd N_x of x in X such that $A \cap N_x \neq \phi$. Then there exists $U \in SO_{\gamma^{*}}(A)$ such that $x \in U \subseteq N_x$. Therefore, $U \cap A = \phi$, so that $A \subseteq X - U$. Clearly $X - U$ is γ^{*}-semi-closed in X and hence $scl_{\gamma^{*}}(A) \subseteq X - U$. Since $x \notin X - U$, we obtain $x \notin scl_{\gamma^{*}}(A)$. This is contradiction to the hypothesis. This proves the necessity.

Conversely, suppose that every γ-semi-nbd of x in X meets A. If $x \notin scl_{\gamma^*}(A)$, then by definition there exists a γ^*-semi-closed F of X such that $A \subseteq F$ and $x \notin F$. Therefore we have $x \in X - F \in SO_{\gamma^*}(X)$. Hence $X - F$ is γ-semi-nbd of x in X. But $(X - F) \cap A = \phi$. This is contradiction to the hypothesis. Thus $x \in scl_{\gamma^*}(A)$.

3 γ-Semi-Open Functions

Theorem 3.1. Let $f : X \to Y$ be a function from a space X into a space Y and γ is an open, monotone and regular operation. Then the following statements are equivalent:
(1) f is γ-semi-open.
(2) $f(int_\gamma(A)) \subseteq sin_{\gamma^*}(f((A))$ for each subset A of X.
(3) For each $x \in X$ and each γ-open-nbd U of x, there exists a γ-semi-nbd V of $f(x)$ such that $V \subseteq f(U)$.

Proof. (1) $\Rightarrow$ (2). Suppose that f is γ-semi-open, and let A be an arbitrary subset of X. Since $f(int_\gamma(A))$ is γ^*-semi-open and $f(int_\gamma(A)) \subseteq f(A)$, then $f(int_\gamma(A)) \subseteq sin_{\gamma^*}(f((A))$.

(2) $\Rightarrow$ (3). Let U be an arbitrary γ-open-nbd of $x \in X$. Then there exists γ-open set O such that $x \in O \subseteq U$. By hypothesis, we have $f(O) = f(int_\gamma(O)) \subseteq sin_{\gamma^*}(f((O))$ and hence $f(O) \subseteq sin_{\gamma^*}(f((O))$. Therefore it follows that $f(O)$ is γ-semi-open-nbd in Y such that $f(x) \in f(O) \subseteq f(U)$. This proves (3).

(3) $\Rightarrow$ (1). Let U be an arbitrary γ-open set in X. For each $y \in f(U)$, by hypothesis there exists a γ-semi-nbd V_y of y in Y such that $V_y \subseteq f(U)$. Since V_y is a γ-semi-nbd of y, there exists a γ^*-semi-open set A_y in Y such that $y \in A_y \subseteq V_y$. Therefore $f(U) = \bigcup\{A_y : y \in f(U)\}$ is a γ^*-semi-open in Y, since is γ regular [9]. This shows that f is a γ-semi-open function.

Theorem 3.2. A bijective function $f : X \to Y$ is γ-semi-open if and only if $f^{-1}(scl_{\gamma^*}(B)) \subseteq cl_\gamma(f^{-1}(B))$ for every subset B of Y, where γ is an open operation.

Proof. Let B be an arbitrary subset of Y. Put

$$U = X - cl_\gamma(f^{-1}(B)) \text{ ...... (I)}$$

Clearly U is a γ-open set in X. Then by hypothesis, $f(U)$ is a γ^*-semi-open set in Y, or $Y - f(U)$ is γ^*-semi-closed set in Y. Since f is onto, from (I), it follows $B \subseteq Y - f(U)$. Thus we have

$scl_{\gamma^*}(B) \subseteq Y - f(U)$. Since f is one-one, we have $f^{-1}(scl_{\gamma^*}(B)) \subseteq f^{-1}(Y) - f^{-1}f(U) = X - f^{-1}f(U) \subseteq X - U = cl_\gamma(f^{-1}(B))$. This proves the necessity.

Conversely, let U be an arbitrary -open set in X. Put $B = Y - f(U)$. Since f is bijective, therefore by hypothesis, $f(U) \cap scl_{\gamma^*}(B) = f(U \cap f^{-1}(scl_{\gamma^*}(B))) \subseteq f(U \cap cl_\gamma(f^{-1}(B)))$. Since U is γ-open, therefore by Lemma 2(3) [14], we have $U \cap cl_\gamma(f^{-1}(B)) \subseteq cl_\gamma(U \cap f^{-1}(B))$. Moreover, it is obvious that $U \cap f^{-1}(B) = \phi$. Thus we have $f(U) \cap scl_{\gamma^*}(B) = \phi$ and hence $scl_{\gamma^*}(B) \subseteq Y - f(U) = B$. Therefore B is a γ^*-semi-closed in Y and hence $f(U)$ is a γ^*-semi-open set in Y. This proves that f is a γ-semi-open mapping.

Definition 3.3 [13]. A function $f : (X, \tau, \gamma) \to (Y, \delta, \beta)$ is said to be (γ, β)-continuous, if for each $x \in X$ and each open set V containing $f(x)$, there exists an open set U such that $x \in U$ and $f(U^\gamma) \subseteq V^\beta$, where γ and β are operations on τ and δ respectively.

Definition 3.4 [13]. A function $f : (X, \tau, \gamma) \to (Y, \delta, \beta)$ is said to be (γ, β)-open (closed), if for any γ-open (closed) set A of X, $f(A)$ is γ-open (closed) in Y.

Theorem 3.5 [1]. Let $f : (X, \tau, \gamma) \to (Y, \delta, \beta)$ be a function and β be an open operation on Y. Then f is (γ, β)-continuous if and only if for each β-open set V in Y, $f^{-1}(V)$ is γ-open in X.

Theorem 3.6 [1]. Let $f : (X, \tau, \gamma) \to (Y, \delta, \beta)$ be a function and β be an open operation on Y. Then the following are equivalent:

(1) f is (γ, β)-open .

(2) $f^{-1}(cl_\beta(B)) \subseteq cl_\gamma(f^{-1}(B))$.

(3) $f^{-1}(bd_\beta(B)) \subseteq bd_\gamma(f^{-1}(B))$ for any subset B of Y.

Theorem 3.7. If a function $f : (X, \tau, \gamma) \to (Y, \delta, \beta)$ is a (γ, β)-open and a (γ, β)-continuous, then the inverse image $f^{-1}(B)$ of each β^*-semi-open set B in Y is γ^*-semi-open in X, where β is an open operation on Y.

Proof. Let B be an arbitrary β^*-semi-open set in Y. Then there exists β-open set V in Y such that $V \subseteq B \subseteq cl_\beta(V)$. Since f is (γ, β)-open, using Theorem 3.6, we have $f^{-1}(V) \subseteq f^{-1}(B) \subseteq$

$f^{-1}(cl_\beta(V)) \subseteq cl_\gamma(f^{-1}(V))$. Since is (γ, β)-continuous and V is β-open in Y, by Theorem 3.5, $f^{-1}(V)$ is γ-open in X. This shows that $f^{-1}(B)$ is γ^*-semi-open set in X.

Theorem 3.8. Let X, Y and Z be three spaces and let $f : X \to Y$ be a function, $g : Y \to Z$ be an injective function and $gof : X \to Z$ is a γ-semi-open function. Then we have:
(1) If f is (γ, β)-continuous and surjective, then g is γ-semi-open.
(2) If g is (β, α)-open, (β, α)-continuous and injective, then f is γ-semi-open, where β is open operation on Y.

Proof. (1) Let V be a β-open set in Y. Then $f^{-1}(V)$ is γ-open in X, because f is (γ, β)-continuous. Since gof is γ-semi-open and f is surjective, therefore $g(V) = (gof)(f^{-1}(V))$ is α^*-semi-open in Z. This shows that g is a γ-semi-open function.
(2) Since g is injective, therefore for $A \subseteq X$, $f(A) = g^{-1}(g(f(A)))$. Let U be a γ-open set in X, then $gof(U)$ is α^*-semi-open. Thus by Theorem 3.7, $g^{-1}(g(f(U))) = f(U)$ is β^*-semi-open in Y. This shows that f is a γ-semi-open function.

Let $B \subseteq X$, $\gamma : \tau \to P(X)$ be an operation. We define $\gamma_B : \tau_B \to P(X)$ as $\gamma_B(U \cap B) = \gamma(U) \cap B$. From here γ_B is an operation and satisfies that $cl_{\gamma_B}(U \cap B) \subseteq cl_\gamma(U \cap B) \subseteq cl_\gamma(U) \cap cl_\gamma(B)$. Using this fact we prove the following:

Theorem 3.9. Let X be a space and B a γ^*-semi-open set in X containing a subset A of X. If A is γ^*-semi-open in the subspace B, then A is γ^*-semi-open in X, where γ is a regular operation.

Proof. Let A be γ_B^*-semi-open in the subspace B. Then there exists a γ_B-open set U_B in B such that $U_B \subseteq A \subseteq cl_{\gamma_B}(U_B)$. Since U_B is γ_B-open in B, there exists a γ-open set U in X such that $U_B = U \cap B$[4]. Thus we have $U \cap B \subseteq A \subseteq cl_{\gamma_B}(U \cap B) \subseteq cl_\gamma(U \cap B) = cl_\gamma(A) \cap cl_\gamma(B)$. Since B is γ^*-semi-open set in X and U is γ-open in X, therefore $U \cap B$ is γ-open in X. Consequently, A is a γ^*-semi-open set in X.

Theorem 3.10. Let X and Y be spaces. If a bijective function $f : X \to Y$ is a γ-semi-open, then for each γ-open set $V(\neq \phi)$ in Y $f|_{f^{-1}(V)} : f^{-1}(V) \to V$ is γ-semi-open, where γ is a regular operation.

Proof. Let U_V be an arbitrary $\gamma_{f^{-1}(V)}$-open set in $f^{-1}(V)$. Then there exists a γ-open set U in X such that $U_V = U \cap f^{-1}(V)$. Now we have $[f|_{f^{-1}(V)}](U_V) = f(U \cap f^{-1}(V)) = f(U) \cap V$. Since $f(U)$ is γ^*-semi-open and V is γ-open, $f(U) \cap V$ is γ^*-semi-open. Hence $[f|_{f^{-1}(V)}](U_V)$ is also γ_V^*-semi-open in V. This shows that $f|_{f^{-1}(V)} : f^{-1}(V) \to V$ is a γ-semi-open mapping.

Theorem 3.11. A bijective function $f : X \to Y$ is γ-semi-open if and only if for any subset V of Y and for any γ-closed set F of X containing $f^{-1}(V)$, there exists a γ^*-semi-closed set G of Y containing V such that $f^{-1}(G) \subseteq F$.

Proof. Let $V \subseteq Y$ and F be a γ-closed set of X containing $f^{-1}(V)$. Put $G = Y - f(X - F)$. Since f is γ-semi-open, so G is γ^*-semi-closed sets in Y . As f is bijective, it follows from $f^{-1}(V) \subseteq F$ that $V \subseteq G$. Calculations give $f^{-1}(G) \subseteq F$.

Conversely, suppose U is γ-open set. Put $V = Y - f(U)$. Then $X - U$ is γ-closed set in X containing $f^{-1}(V)$. By hypothesis, there exists a γ^*-semi-closed set G of Y such that $V \subseteq G$ and $f^{-1}(G) \subseteq (X - U)$. On the other hand, it follows from $V \subseteq G$ that $f(U) = (Y - V) \subseteq (Y - G)$. Therefore, we obtain $f(U) = (Y - G) \in SO_{\gamma^*}(Y)$. This shows that f is γ-semi-open.

Lemma 3.12 [5]. The following properties of a subset A of X are equivalent:

(1) A is γ^*-semi-closed.

(2) $int_\gamma(cl_\gamma(A)) \subseteq A$.

(3) $X - A$ is γ^*-semi-open.

Theorem 3.13. If $f : X \to Y$ is (γ, β)-open and (γ, β)-continuous mapping, then the inverse image $f^{-1}(B)$ of each β^*-semi-closed B in Y is γ^*-semi-closed in X, where β is an open operation on Y.

Proof. This follows from Theorem 3.7 and Lemma 3.12.

Theorem 3.14. Let $f : X \to Y$ be surjective and $g : Y \to Z$ be an injective function and let $gof : X \to Z$ be a γ-semi-closed function. Then

(1) If f is (γ, β)-continuous and surjective, then g is β-semi-closed.

(2) If g is (β, α)-open, (β, α)-continuous and injective, then f is γ-semi-closed, where β is an open

operation on Y.

Proof. (1) Suppose H is an arbitrary β-closed set in Y. Then $f^{-1}(H)$ is γ-closed in X because f is (γ,β)-continuous. Since gof is γ-semi-closed and f is surjective, $gof(f^{-1}(H)) \subseteq g(f(f^{-1}(H))) = g(H)$, is α^*-semi-closed in Z. This implies that g is β-semi-open function. This proves (1).

(2) Since g is injective so for every subset A of X, $f(A) = g^{-1}(g(f(A)))$. Let F be an arbitrary γ-closed set in X. Then $gof(F)$ is γ^*-semi-closed. It follows immediately from Theorem 3.13 that $f(F)$ is γ^*-semi-closed set in Y. This implies that f is γ-semi-closed.

4 γ-Semi-Closed Functions

Theorem 4.1. Let γ be an open and monotone operation. A function $f : X \to Y$ is γ-semi-closed if and only if $f(cl_\gamma(A)) \supseteq int_\gamma(cl_\gamma(f(A)))$ for every subset A of X.

Proof. Suppose f is a γ-semi-closed mapping and A is an arbitrary subset of X. Then $f(cl_\gamma(A))$ is γ^*-semi-closed in Y. Then by Lemma 3.12, we obtain $f(cl_\gamma(A)) \supseteq int_\gamma(cl_\gamma(f(cl_\gamma(A)))) \supseteq int_\gamma(cl_\gamma(f(A)))$. This implies that $f(cl_\gamma(A)) \supseteq int_\gamma(cl_\gamma(f(A)))$.

Conversely, suppose that F is an arbitrary γ-closed set in X. Then by hypothesis, we have $int_\gamma(cl_\gamma(f(F))) \subseteq f(cl_\gamma(F)) = f(F)$. By Lemma 3.12, $f(F)$ is γ^*-semi-closed in Y. This implies that f is γ-semi-closed.

Recall [9] that the intersection of all γ^*-semi-closed sets containing A is called γ-semi-closure of A and is denoted by $scl_{\gamma^*}(A)$. Clearly A is γ^*-semi-closed if and only if $scl_{\gamma^*}(A) = A$.

Theorem 4.2. Let γ be an open and monotone operation. A function $f : X \to Y$ is γ-semi-closed if and only if $scl_{\gamma^*}(A) \subseteq f(cl_\gamma(A))$ for every subset A of X.

Proof. Suppose f is a γ-semi-closed mapping and A is an arbitrary subset of X. Then $f(cl_\gamma(A))$ is γ^*-semi-closed. Since $f(A) \subseteq f(cl_\gamma(A))$, we obtain $scl_{\gamma^*}(f(A)) \subseteq f(cl_\gamma(A))$. This implies $scl_{\gamma^*}(f(A)) \subseteq f(cl_\gamma(A))$.

Sufficiency follows from Theorem 4.1.

Theorem 4.3. A surjective function $f : X \to Y$ is γ-semi-closed if and only if for each subset B in Y and each γ-open set U in X containing $f^{-1}(B)$, there exists a γ^*-semi-open set V in Y containing B such that $f^{-1}(V) \subseteq U$, where γ is a monotone and regular operation.

Proof. Suppose B is an arbitrary subset in Y and U is an arbitrary γ-open set in X containing $f^{-1}(B)$. We put

$$V = Y - f(X - U) \text{ ..... (*)}$$

Then V is γ^*-semi-open set in Y. Since $f^{-1}(B) \subseteq U$, calculations give $B \subseteq V$. Moreover, by (*), we have $f^{-1}(V) = f^{-1}(Y) - f^{-1}(f(X-U)) = X - f^{-1}(f(X-U)) \subseteq X - (X-U) = U$.

Conversely, suppose that F is an arbitrary γ-closed set in X. Let y be an arbitrary point in $Y - f(F)$, then $f^{-1}(y) \subseteq X - f^{-1}(f(F)) \subseteq X - F$, and $X - F$ is γ-open in X. Hence by the hypothesis, there exists a γ^*-semi-open set V_y containing y such that $f^{-1}(V_y) \subseteq X - F$. This implies that $v \in V_y \subseteq Y - f(F)$. We obtain that $Y - f(F) = \bigcup\{V_y : y \in Y - f(F)\}$ is γ^*-semi-open in Y, since union of any collection of γ^*-semi-open sets is γ^*-semi-open. Therefore $f(F)$ is γ^*-semi-closed.

5 γ-Semi-Continuous Functions

Theorem 5.1. Let $f : X \to Y$ be a function and γ is an open operation. Then the following are equivalent:

(1) f is γ-semi-continuous.

(2) $int_\gamma(cl_\gamma(f^{-1}(B))) \subseteq f^{-1}(cl_\gamma(B))$ for each subset B of Y.

(3) $f(int_\gamma(cl_\gamma(A))) \subseteq cl_\gamma(f(A))$ for each subset A of X.

Proof. (1) ⇒ (2). Let B be an arbitrary subset of Y. Then by (1), $f^{-1}(cl_\gamma(B))$ is a γ^*-semi-closed set of X. Since $B \subseteq cl_\gamma(B)$, by Lemma 3.12, we get $f^{-1}(cl_\gamma(B)) \supseteq int_\gamma(cl_\gamma(f^{-1}(cl_\gamma(B)))) \supseteq int_\gamma(cl_\gamma(f^{-1}(B)))$implies that $int_\gamma(cl_\gamma(f^{-1}(B))) \subseteq f^{-1}(cl_\gamma(B))$.

(2) ⇒ (3). Let A be an arbitrary subset of X. Put $B = f(A)$. Then $A \subseteq f^{-1}(B)$. Therefore by hypothesis, we have $int_\gamma(cl_\gamma(A)) \subseteq int_\gamma(cl_\gamma(f^{-1}(B))) \subseteq f^{-1}(cl_\gamma(B))$. Consequently, we have $f(int_\gamma(cl_\gamma(A))) \subseteq ff^{-1}(cl_\gamma(B)) \subseteq cl_\gamma(B) = cl_\gamma(f(A))$. This gives (3).

(3) ⇒ (1). Let F be an arbitrary γ-closed set of Y. Put $A = f^{-1}(F)$, then $f(A) \subseteq F$. Therefore by hypothesis, we have

$$f(int_\gamma(cl_\gamma(A))) \subseteq cl_\gamma(f(A)) \subseteq cl_\gamma(F) = F \text{ ..... (**)}$$

By (**), we have $int_\gamma(cl_\gamma(A)) \subseteq f^{-1}f(int_\gamma(cl_\gamma(A))) \subseteq f^{-1}(cl_\gamma(f(A))) \subseteq f^{-1}(cl_\gamma(F)) = f^{-1}(F)$, or $int_\gamma(cl_\gamma(A)) \subseteq f^{-1}(F)$,. By Lemma 3.12, $f^{-1}(F)$ is a γ^*-semi-closed set in X. This implies that f is γ-semi-continuous.

Definition 5.2. Let X be a space $A \subseteq X$ and $p \in X$. Then p is a γ^*-semi-limit point of A, for all γ^*-semi-open set U containing p such that $U \cap (A - \{p\}) \neq \phi$. The set of all γ^*-semi-limit point of A is said to be γ^*-semi-derived set of A and is denoted by $sd_{\gamma^*}(A)$.

Clearly if $A \subseteq B$ then $sd_{\gamma^*}(A) \subseteq sd_{\gamma^*}(B)$..... (I)

Remark 5.3. From the definition, it follows that p is a γ^*-semi-limit point of A if and only if $p \in scl_{\gamma^*}(A - \{p\})$.

Theorem 5.4. The γ^*-semi-derived set, sd_{γ^*}, has the following properties:
(1) $scl_{\gamma^*}(A) = A \cup sd_{\gamma^*}(A)$.
(2) $sd_{\gamma^*}(A \cup B) = sd_{\gamma^*}(A) \cup sd_{\gamma^*}(B)$. In general
(3) $\bigcup_i sd_{\gamma^*}(A_i) = sd_{\gamma^*}(\bigcup_i (A_i))$.
(4) $sd_{\gamma^*}(sd_{\gamma^*}(A)) \subseteq sd_{\gamma^*}(A)$.
(5) $scl_{\gamma^*}(sd_{\gamma^*}(A)) = sd_{\gamma^*}(A)$.

Proof. (1) Let $x \in scl_{\gamma^*}(A)$. Then $x \in C$, for every γ^*-semi-closed superset C of A. Now
(i) If $x \in A$, then $x \in A \cup sd_{\gamma^*}(A)$.
(ii) If $x \notin A$, then we prove that $x \in scl_{\gamma^*}(A)$.
To prove (ii), suppose U is γ^*-semi-open set containing x. Then $U \cap A \neq \phi$, for otherwise, $A \subseteq X - U = C$, where C is a γ^*-semi-closed superset of A not containing x. This contradicts the fact that x belongs to every γ^*-semi-closed superset C of A. Therefore $x \in sd_{\gamma^*}(A)$ gives $x \in A \cup sd_{\gamma^*}(A)$.

Conversely, suppose that $x \in A \cup sd_{\gamma^*}(A)$, we show that $x \in scl_{\gamma^*}(A)$. If $x \in A$ then $x \in scl_{\gamma^*}(A)$. If $x \in sd_{\gamma^*}(A)$, then we show that x is in every γ^*-semi-closed superset of A. We suppose otherwise that there is γ^*-semi-closed superset C of A not containing x. Then $x \in X - C = U$(say), which is γ^*-semi-open and $U \cap A = \phi$. This implies that $x \notin sd_{\gamma^*}(A)$. This contradiction proves that $x \in scl_{\gamma^*}(A)$. Consequently $scl_{\gamma^*}(A) = A \cup sd_{\gamma^*}(A)$. This proves (1).

(2) $sd_{\gamma^*}(A \cup B) \subseteq sd_{\gamma^*}(A) \cup sd_{\gamma^*}(B)$.

Let $x \in sd_{\gamma^*}(A \cup B)$. Then $x \in scl_{\gamma^*}((A \cup B) - \{x\})$ or $x \in scl_{\gamma^*}((A - \{x\}) \cup (B - \{x\})$ implies $x \in scl_{\gamma^*}(A - \{x\})$ or $x \in scl_{\gamma^*}(B - \{x\})$. This gives $x \in sd_{\gamma^*}(A)$ or $x \in sd_{\gamma^*}(B)$. Therefore $x \in sd_{\gamma^*}(A) \cup sd_{\gamma^*}(B)$. This proves $sd_{\gamma^*}(A \cup B) \subseteq sd_{\gamma^*}(A) \cup sd_{\gamma^*}(B)$

Converse follows directly by using the property(I).

(3) The proof is immediate by property (I).

(4) Suppose that $x \notin sd_{\gamma^*}(A)$. Then $x \notin scl_{\gamma^*}(A - \{x\})$. This implies that there is γ^*-semi-open set U such that $x \in U$ and $U \cap (A - \{x\}) = \phi$. We prove that $x \notin sd_{\gamma^*}(sd_{\gamma^*}(A))$. Suppose on the contrary that $x \in sd_{\gamma^*}(sd_{\gamma^*}(A))$. Then $x \in scl_{\gamma^*}(sd_{\gamma^*}(A) - \{x\})$. Since $x \in U$, we have $U \cap (sd_{\gamma^*}(A) - \{x\}) \neq \phi$. Therefore there is a $q \neq x$ such that $q \in U \cap (sd_{\gamma^*}(A))$. It follows that $q \in (U - \{x\}) \cap (sd_{\gamma^*}(A) - \{x\})$. Hence $(U - \{x\}) \cap (sd_{\gamma^*}(A) - \{x\}) \neq \phi$, a contradiction to the fact that $(U \cap (sd_{\gamma^*}(A) - \{x\}) = \phi$. This implies that $x \notin sd_{\gamma^*}(sd_{\gamma^*}(A))$ and so $sd_{\gamma^*}(sd_{\gamma^*}(A)) \subseteq sd_{\gamma^*}(A)$. This proves (4).

(5) This is a consequence of (1), (2) and (4).

Theorem 5.5 [5]. Let $f : X \to Y$ be a function. Then the following are equivalent:
(1) $f : X \to Y$ is γ-semi-continuous.
(2) $scl_{\gamma^*}(f^{-1}(A)) \subseteq f^{-1}(cl_{\gamma}(A))$ for each subset A of Y.

Theorem 5.6. Let $f : X \to Y$ be a function and γ is an open operation. Then the following are equivalent:
(1) $f : X \to Y$ is γ-semi-continuous.
(2) $f(sd_{\gamma^*}(A)) \subseteq cl_{\gamma}(f(A))$ for any subset A of X.

Proof. (1) $\Rightarrow$ (2). Suppose that f is γ-semi-continuous. Let A be any set in X. Since $cl_{\gamma}(f(A))$ is γ-closed in Y. $f^{-1}(cl_{\gamma}(A))$ is γ^*-semi-closed in X. $A \subseteq f^{-1}(f(A)) \subseteq f^{-1}(cl_{\gamma}(f(A)))$ gives $scl_{\gamma^*}(A) \subseteq scl_{\gamma^*}(f^{-1}(cl_{\gamma}(f(A)))) = f^{-1}(cl_{\gamma}(f(A)))$. Therefore $f(sd_{\gamma^*}(A)) \subseteq f(scl_{\gamma^*}(A)) \subseteq ff^{-1}(cl_{\gamma}(f(A))) \subseteq cl_{\gamma}(f(A))$. Consequently, $f(sd_{\gamma^*}(A)) \subseteq cl_{\gamma}(f(A))$.

(2) $\Rightarrow$ (1). Suppose that $f(sd_{\gamma^*}(A)) \subseteq cl_{\gamma}(f(A))$, for $A \subseteq X$. Let B be any γ-closed subset of Y. We show that $f^{-1}(B)$ is γ^*-semi-closed in X. By hypothesis, $f(sd_{\gamma^*}(f^{-1}(B))) \subseteq cl_{\gamma}(f(f^{-1}(B))) \subseteq cl_{\gamma}(B) = B$ or $f(sd_{\gamma^*}(f^{-1}(B))) \subseteq B$ gives $sd_{\gamma^*}(f^{-1}(B)) \subseteq f^{-1}f(sd_{\gamma^*}(f^{-1}(B))) \subseteq f^{-1}(B)$ or $sd_{\gamma^*}(f^{-1}(B)) \subseteq f^{-1}(B)$ implies $f^{-1}(B)$ is γ^*-semi-closed in X. Thus f is γ-semi-continuous.

Theorem 5.7 [5]. Let $f : X \to Y$ be a function and $x \in X$. Then f is γ-semi-continuous if and only if for each γ-open set B containing f(x), there exists $A \in SO_{\gamma^*}(X)$ such that $x \in A$ and $f(A) \subseteq B$, where γ is a regular operation.

We use Theorem 5.7 and prove the following:

Theorem 5.8. Let $f : X \to Y$ be an injective function. If f is γ-semi-continuous then $f(sd_{\gamma^*}(A)) \subseteq (f(A))^d_\gamma$ for every $A \subseteq X$, where γ is a regular operation.

Proof. Suppose that f is γ-semi-continuous. Let $A \subseteq X$, $a \in sd_{\gamma^*}(X)$ and V be a γ-open-nbd of $f(a)$. Since f is γ-semi-continuous then by Theorem 5.7, there exists a γ-semi-open-nbd U of a such that $f(U) \subseteq V$. But $a \in sd_{\gamma^*}(A)$, therefore there exists an element $a_1 \in U \cap A$ such that $a \neq a_1$; then $f(a_1) \in f(A)$ and since f is an injection $f(a) \neq f(a_1)$. Thus every γ-open-nbd V of $f(a)$ contains an element $f(a_1)$ of $f(A)$ different from $f(a)$. Consequently $f(a) \in (f(A))^d_\gamma$. We have therefore, $f(sd_{\gamma^*}(A)) \subseteq (f(A))^d_\gamma$.

The following theorem follows from Theorem 5.6:

Theorem 5.9. Let $f : X \to Y$ be a function. If for every $A \subseteq X$, $f(sd_{\gamma^*}(A)) \subseteq (f(A))^d_\gamma$, then f is γ-semi-continuous, where γ is an open operation.

Theorem 5.10. A function $f : X \to Y$ is γ-semi-continuous if and only if $f^{-1}(int_\gamma(B)) \subseteq sint_{\gamma^*}(f^{-1}(B))))$, for each $B \subseteq Y$, where γ is a regular operation.

Proof. For any $B \subseteq Y$, $int_\gamma(B) = Y - cl_\gamma(Y - B)$ [14]. This implies $f^{-1}(int_\gamma(B)) = f^{-1}(Y - cl_\gamma(Y - B)) = X - f^{-1}(cl_\gamma(Y - B))$. Since f is γ-semi-continuous, by Theorem 5.5 we have $scl_{\gamma^*}(f^{-1}(Y - B)) \subseteq f^{-1}(cl_\gamma(Y - B))$. Hence $f^{-1}(int_\gamma(B)) \subseteq X - scl_{\gamma^*}(f^{-1}(Y - B))$. Thus $f^{-1}(int_\gamma(B)) \subseteq X - scl_{\gamma^*}(X - f^{-1}(B))$. Hence $f^{-1}(int_\gamma(B)) \subseteq X - scl_{\gamma^*}(X - f^{-1}(B)) = sint_{\gamma^*}(f^{-1}(B))))$.

Conversely, let B be an arbitrary γ-open set in Y. Then $int_\gamma(B) = B$. By hypothesis $f^{-1}(B) = f^{-1}(int_\gamma(B)) \subseteq sint_{\gamma^*}(f^{-1}(B))$ implies $f^{-1}(B) \subseteq sint_{\gamma^*}(f^{-1}(B))$. But $sint_{\gamma^*}(f^{-1}(B)) \subseteq f^{-1}(B)$. Therefore, $f^{-1}(B) = sint_{\gamma^*}(f^{-1}(B))$. Thus $f^{-1}(B)$ is γ^*-semi-open . Consequently, f is γ-semi-continuous.

References

[1] B. Ahmad and F. U. Rehman: *Operations on Topological Spaces-II, Math. Today,* 11 (1993),13-20.

[2] B. Ahmad and S. Hussain: *Properties of γ-Operations on Topological Spaces, Aligarh Bull. Math.,* 22 (1) (2003),45-51.

[3] B. Ahmad and S. Hussain: *γ- Convergence in Topological Spaces, Southeast Asian Bull. Math.,* 29 (5)(2005), 835-842.

[4] B. Ahmad and S. Hussain: *γ^*- regular and γ-normal Spaces, Math. Today,* 22 (1)(2006), 37-44.

[5] B. Ahmad and S. Hussain: *γ^*-Semi-Open Sets in Topological Spaces, Southeast Asian Bull. Math.,* 34(6)(2010),(In Press).

[6] C. Carpintero, E. Rosas, J. Vielma: *Operadores Asociados a uns Topologia Γ Sobre un Conjunto X y Nociones Conexas, Divulgaciones Mathematicas,* 6 (2)(1998), 139-144.

[7] A. Csaszar : *Generalized Open Sets, Acta Math. Hungar.,* 75(1997), 65-87.

[8] A. Csaszar : *Generalized Topology, Generalized Continuity, Acta Math. Hungar.,* 96(2002),351-357.

[9] S. Hussain, B. Ahmad and T. Noiri : *γ^*-Semi-Open Sets in Topological Spaces , Asian Eur. J. Math.,* 3(3)(2010), 427-433. DOI: 10.1142/S1793557110000337.

[10] S. Kasahara: *Operation- Compact Spaces, Math. Japon.,* 24(1979), 97-105.

[11] N. Levine: *Semi-Open Sets and Semi Continuity in Topological Spaces, Amer. Math. Monthly,* 70(1963), 36-41.

[12] S. N. Meheshwari and R. Prasad: *On R_0-spaces, Portugal Math.,* 34(1975), 213-217.

[13] H. Ogata : *Operations on Topological Spaces and Associated Topology , Math. Japon.,* 36 (1) (1991), 175-184.

[14] F.U. Rehman and B. Ahmad: *Operations on Topological Spaces-I, Math. Today,* 10(1992), 29-36.